



Supplementary Table1. List of primers used in q-PCR

Gene	Sense (5'-3')	Antisense (5'-3')
<i>Lbp</i>	TCGTGGGCAGTACGAGTTTC	AAGAGATTCAGCAGCCACCC
<i>Cd14</i>	GTCAGGAACTCTGGCTTTGC	TGGCTTTTACCCACTGAACC
<i>Srebf1</i>	GATCAAAGAGGAGCCAGTGC	TAGATGGTGGCTGCTGAGTG
<i>Fas</i>	TGGGTTCTAGCCAGCAGAGT	ACCACCAGAGACCGTTATGC
<i>Acc1</i>	GCCTCTTCCTGACAAACGAG	TGACTGCCGAAACATCTCTG
<i>Ppara</i>	GAGGGTTGAGCTCAGTCAGG	GGTCACCTACGAGTGGCATT
<i>Pparg</i>	CTGTGAGACCAACAGCCTGA	AATGCGAGTGGTCTTCCATC
<i>Tnfa</i>	TTCTATGGCCCAGACCCTCA	TGGTTTGCTACGACGTGGG
<i>Il1b</i>	GCCCATCCTCTGTGACTCAT	AGGCCACAGGTATTTTGTCTG
<i>Il6</i>	AGTTGCCTTCTTGGGACTGA	TCCACGATTTCCAGAGAAC
<i>Acta2</i>	CTGACAGAGGCACCACTGAA	CATCTCCAGAGTCCAGCACA
<i>Col1a1</i>	GAGCGGAGAGTACTGGATCG	GCTTCTTTTCCTTGGGGTTC
<i>Tgfb1</i>	TTGCTTCAGCTCCACAGA GA	TGGTTGTAGAGGGCAAGGAC
<i>Gapdh</i>	CCACCCAGAAGACTGTGGAT	CACATTGGGGGTAGGAACAC